

A DFT study of (001) CaSnO_3 orthorhombic perovskite surface with adsorbed gases (NH_3 and CO)

Jefferson Maul¹, I. M. G. Santos² and J. R. Sambrano³

¹Dipartimento di Chimica, Università degli studi di Torino, Via Giuria 5, 10125 Torino, Italy

²Núcleo de Pesquisa e Extensão: Laboratório de Combustíveis e Materiais, Universidade Federal da Paraíba, CEP 58051-900, João Pessoa, PB, Brazil

³Computational Simulation Group, CDMF-UNESP, São Paulo State University, CEP 17033-360, Bauru, SP, Brazil
email:jmauldea@unito.it

The CaSnO_3 (001) surface is here studied under the Density Functional Theory (DFT) framework with PBE and PBE0 functionals using Crystal14 program[1]. Two terminations of the pristine surface were studied, the first was surface terminated in CaO (CaO-T) that presented to be slightly more stable than the SnO_2 terminated, $\text{SnO}_2\text{-T}$. Carbon monoxide, CO , and ammonia and NH_3 gases were adsorbed over the $\text{SnO}_2\text{-T}$ surface, said to be more reactive. The surfaces and adsorptions were evaluated using differential charge density maps, Density of States (DOS) and Raman intensities (through the Coupled-Perturbed-Hartree-Fock/Kohn-Sham algorithm). The CO adsorption was weaker than the NH_3 one. Both presented large shifts on the Raman spectra (Figs. 1 and 2), however only NH_3 presented a new intense Raman peak about 3098 cm^{-1} , this assigned to the stretch of the the hydrogen towards the superficial oxygen. The (001) of the CaSnO_3 perovskite presented sensor properties for the studied gases, and its interaction with NH_3 is noteworthy.




Fig1: Raman spectra of CO adsorption on $\text{SnO}_2\text{-T}$ at PBE0 level




Fig2: Raman spectra of NH_3 adsorption on $\text{SnO}_2\text{-T}$ at PBE0 level

References:

[1] R. Dovesi, V. R. Saunders, C. Roetti et al., CRYSTAL14 User's Manual (2014).