



3rd Science for the Environment Conference
Aarhus Denmark 1-2 October 2015

SURVEY EFFORT ALLOCATION USING ADVANCED DESIGN: CLAM POPULATION AS CASE STUDY

Nathalie Caill-Milly*, Frank D'Amico**, Noëlle Bru**, Florence Sanchez*,
Jennifer Brown***, Muriel Lissardy*

* Ifremer, Laboratory Halieutic Resources from Aquitaine, FED 4155 – MIRA, France

** UPPA, Laboratory of Mathematics and their Applications of Pau – UMR CNRS 5142, FED

4155 – MIRA, France. *** University of Canterbury, Department of Mathematics and Statistics,
New Zealand

ABSTRACT

The planning strategies for the Manila clam (*Ruditapes philippinarum*) resource in Arcachon Bay use an adaptive co-management approach, linking scientific advice with fishermen's current practice. This management framework is based on data collected from field surveys. Surveys started in 2000 and have continued biennially since 2006. Here we describe work to assess and compare the performances of different sample design options for monitoring the stock of this infaunal bivalve, using simulation methodology. Random stratified sampling (RSS) is a sampling protocol used most commonly for surveys of this nature, and at this scale. We compared RSS with recently developed, spatially balanced survey designs. One of these designs is generalized random tessellation stratified (GRTS) which has mainly been used for aquatic resources characterized by linear networks. Here we extend its application to fisheries. In our research the steps followed were: 1) creation of a virtual population using geostatistics based on the existing data, 2) selection of sampling stations (optimal number and locations) with RSS and GRTS designs using statistical resampling methods, 3) interpolation of the biomass and abundance values relating to the selected stations to compute estimations, 4) comparison of the designs' precision and costs. In our study GRTS performed better than RSS for estimating survey parameters. These results are encouraging but need to be extended for other surveys with different sized strata. Integration of the operating costs complements our findings. GRTS emerges as promising sample design both in terms of optimizing the design and satisfying financial constraints. The practical implementation of the design into large-scale operational surveys is discussed. The design allows the large-scale survey to be retained even in an environment of decreased funding for monitoring. The GRTS design will be a useful design for other exploited bivalves' populations that are important components of the marine benthic ecosystem.

References

Benedetti, R, Piersimoni, F, Postiglione, P., 2014. Sampling spatial units for agricultural surveys. Advances in Spatial Science Series. Springer.

Caill-Milly N., 2012. Relations entre l'état d'une ressource et son exploitation via la compréhension et la formalisation des interactions de socio-écosystèmes. Application à la palourde japonaise (*Venerupis philippinarum*) du bassin d'Arcachon. PhD Thesis, Université



**3rd Science for the Environment Conference
Aarhus Denmark 1-2 October 2015**

de Pau et des Pays de l'Adour (UPPA).

Muggeo, V.M.R., Adelfio, G., 2011. Efficient change point detection in genomic sequences of continuous measurements. *Bioinformatics*. 27: 161-166.

Stevens, D. L., Jr. and A. R., Olsen, 2004. Spatially-balanced sampling of natural resources. *Journal of American Statistical Association*. 99(465): 262-278.

