

The role of PAHs in the formation of H₂ in the interstellar medium

R. Jaganathan¹, D. Campisi², J. Palotas³, A. Cassidy¹, L. Hornekær¹

¹ *Department of Physics and Astronomy, Aarhus University, Aarhus, Denmark*

² *Faculty of Science, Leiden University, Leiden, The Netherlands*

³ *FELIX Laboratory, Radboud University, Nijmegen, The Netherlands*


rijutha@phys.au.dk

Carbonaceous surfaces are known to act as catalysts for the formation of molecular hydrogen in the interstellar medium [1,2]. Molecular hydrogen is the most abundant molecule in the universe and it controls the chemistry of the interstellar medium. Hence, the formation of molecular hydrogen is the first step in the evolution of the chemical complexity of the interstellar medium.

Observations correlate the abundance of polycyclic aromatic hydrocarbons (PAHs) to an increased rate of H₂ formation [3]. Experiments and theoretical calculations indicate that PAHs play a role as catalysts for H₂ formation [4]. Here we investigate the role that pentacene (C₂₂H₁₄), a model PAH, might play in reactions with atomic H. Pentacene, a linear PAH consisting of five fused benzene rings, is also commonly used to make organic electronic devices like thin film field effect transistors [5].

A monolayer of pentacene was prepared under ultra-high vacuum conditions on a Au (111) surface and then exposed to a controlled fluence of hydrogen atoms. X-ray photoelectron spectroscopy was used to characterize the system by tracking chemical shifts in the C1s and Au4f core levels.

After hydrogenation, a chemical shift was observed for the carbon core level electrons. This indicates that the sp² carbon sites change to a sp³ hybridisation after hydrogen exposure. Furthermore, there is evidence for more than one type of sp³ hybridised carbon site. Chemical shifts in the gold core levels were observed firstly after dosing of pentacene and again after hydrogen exposure. The latter could indicate Au-C bond formation simultaneous with new C-H formation on the pentacene molecule or a Au-H interaction, or both. Combining this data with STM images and DFT calculations will help us understand the dynamics of hydrogen addition to the pentacene molecule and elucidate a route towards superhydrogenation of the PAHs in the interstellar medium.



References:

- [1] V. Pirronello, C. Liu, et al., *Astron. Astrophys.* **344**, 681–686 (1999).
- [2] I. Alata, G. A. Cruz-Diaz, et al., *Astron. Astrophys.* **569**, A119 (2014).
- [3] E. Habart, F. Boulanger, et al., *Astron. Astrophys.* **397**, 623-634 (2003).
- [4] V. Mennella, L. Hornekær, et al., *The Astrophysical Journal Letters*, **745**, L2 (2012).
- [5] M. Lu, R. Wang, et al., *Phys.: Condens. Matter* **28**, 094005 (2016).